

## Are We Ready for Mass Screening to Detect Atrial Fibrillation?

Jeffrey S. Healey, MD; Roopinder K. Sandhu, MD

Atrial fibrillation (AF) is the most common cardiac arrhythmia<sup>1</sup> and a leading cause of stroke.<sup>2</sup> AF-related strokes are associated with significant morbidity, mortality, and healthcare costs,<sup>2-5</sup> yet they are highly preventable.<sup>6,7</sup> Unfortunately, AF is often undiagnosed or untreated when stroke occurs.<sup>8</sup> Given the availability of effective oral anticoagulant (OAC) medications and evidence-based guidelines for their use,<sup>9-11</sup> population-based AF screening has the potential to become an important public health program.

---

### Article see p 2176

---

There is considerable interest in developing AF screening programs for 3 reasons: (1) The prevalence of AF is increasing as a result of our aging population; (2) new OAC medications offer safe, effective, and convenient therapy for patients once AF is identified; and (3) a variety of portable, wearable, and implantable technologies have been developed to detect AF, which may facilitate AF screening in a variety of clinical settings.<sup>12-14</sup> Although there is great optimism for AF screening, high-quality studies examining optimal AF screening methods, settings, and target populations are only starting to emerge.<sup>12,13,15</sup>

In this issue of *Circulation*, Svennberg and colleagues<sup>16</sup> report the results of the STROKESTOP study, a prospective, population-based study of systematic AF screening with intermittent ambulatory ECG recordings among those 75 and 76 years of age in Stockholm County and the Halland region in Sweden. Of the 14 387 individuals invited, an impressive 54% (7173) participated and transmitted 30-second, single-lead ECG recordings (average, 26.4 per participant) over a 2-week period.<sup>16</sup> AF incidence was 3.0%, which is higher than the 1.4% observed in a systematic review of studies using single-time-point screening in a range of populations, some of which included patients as young as 18 years of age.<sup>15</sup> An earlier study by this group, which had a 5.2% rate of incident AF, also found that only 25% of incident cases were diagnosed with the baseline 12-lead ECG, with the remainder detected during 2 weeks of twice-daily home ECG monitoring.<sup>17</sup> Compared with single-time-point screening, which typically detects permanent or persistent AF, the use of multiple ECG

transmissions over time facilitates the detection of paroxysmal AF, thereby increasing the rate of AF detection. This is consistent with prior studies of prolonged AF monitoring in other high-risk populations.<sup>18,19</sup>

In the STROKESTOP study, 9.3% of patients screened reported a previous diagnosis of AF, and 77.6% of these patients were receiving OAC therapy at the time of screening.<sup>16</sup> This represents a much higher background rate of OAC use among AF patients than has previously been reported in real-world settings,<sup>20,21</sup> possibly reflecting better guideline adherence and patient acceptance of OAC than elsewhere. Among the participants in the STROKESTOP study with newly diagnosed AF, OAC therapy was started in 93%, with almost three quarters receiving a new OAC medication.<sup>16</sup> Among patients with known but untreated AF, OAC therapy was started in 47% of patients after identification of AF on screening. Of patients who did not start OAC, only 14% had contraindications; the main reasons for not starting an OAC medication were patient or physician preference.<sup>16</sup>

In STROKESTOP,<sup>16</sup> the very high rates of subject participation and acceptance of OAC allow the authors to draw very accurate conclusions about the true prevalence of undiagnosed AF in this population and the potential to increase the uptake of OAC therapy. However, these study features also make it difficult to extrapolate the findings of STROKESTOP to other settings and countries. The effectiveness of AF screening in other regions will depend on the prevalence of AF, which is influenced by ethnic heterogeneity and differences in mean population age. There is also likely to be regional differences in the feasibility of conducting screening and the ability of local systems to effectively deliver OAC therapy if AF is detected. For example, in an Australian study of 1000 pharmacy customers >65 years of age, pharmacists performed a manual pulse check and an iPhone single-lead ECG.<sup>13</sup> This study found incident AF in 1.5% of individuals, of whom 60% started OAC therapy. Although this experience demonstrated a lower use of OAC than in STROKESTOP, it did demonstrate that screening was both feasible and cost-effective.<sup>13</sup>

The STROKESTOP screening program estimated a cost of €4164 per quality-adjusted life-year in this 75- and 76-year-old population.<sup>16</sup> A similar value was calculated in an earlier study that conducted a modeled cost-effective analysis, extending a pharmacy-based iPhone ECG screening program to the general community in those 65 to 84 years of age.<sup>13</sup> This earlier study estimated an incremental cost-effectiveness ratio of €3142 per quality-adjusted life-year with warfarin and €6267 per quality-adjusted life-year with new OAC therapy.<sup>13</sup> In both studies, the precise cost-effectiveness ratio was sensitive to the rate of participation in screening and the proportion of affected individuals who started and were maintained on OAC.

---

The opinions expressed in this article are not necessarily those of the editors or of the American Heart Association.

From Population Health Research Institute, McMaster University, Hamilton, Ontario, Canada (J.S.H); and University of Alberta, Edmonton, Canada (R.K.S.).

Correspondence to Jeffrey S. Healey, MD, McMaster University, 1280 Main St W, Hamilton, ON L8S4L8, Canada. E-mail jeff.healey@phri.ca (*Circulation*. 2015;131:2167-2168.

DOI: 10.1161/CIRCULATIONAHA.115.017288.)

© 2015 American Heart Association, Inc.

*Circulation* is available at <http://circ.ahajournals.org>

DOI: 10.1161/CIRCULATIONAHA.115.017288

The success and cost-effectiveness of a particular AF screening program will depend not only on the population and screening method but also on the setting of the program. One strength of screening in a pharmacy or family practice setting is that it capitalizes on existing healthcare infrastructure that may allow more sustainability over the long term. Furthermore, AF screening can be combined with existing cardiovascular screening initiatives such as the diagnosis and treatment of hypertension and diabetes mellitus. This may not only save time and increase patient acceptance but also facilitate the delivery of synergistic stroke-prevention therapies such as blood pressure lowering and OAC.

There is little doubt that available screening tools can diagnose AF. However, for large-scale screening to succeed, screening must detect AF and ultimately prevent stroke in a cost-effective fashion. This means screening only populations with sufficient risk of AF and in a setting where screening can be performed accurately and efficiently, preferably in concert with other screening programs. The STROKESTOP study has shown us what is possible under ideal circumstances. More research is needed to determine how well other strategies will work in other settings.

### Sources of Funding

Dr Healey received a personnel award from the Heart and Stroke Foundation, Ontario Provincial office (MC7450).

### Disclosures

Dr Healey has received research grants and speaking fees from Boehringer-Ingelheim, Bayer, Bristol-Meyers-Squibb, Pfizer, Medtronic, and St. Jude Medical and is the medical director of m-Health Solutions. Dr Sandhu reports no conflicts.

### References

- Miyasaka Y, Barnes ME, Gersh BJ, Cha SS, Bailey KR, Abhayaratna WP, Seward JB, Tsang TS. Secular trends in incidence of atrial fibrillation in Olmsted County, Minnesota, 1980 to 2000, and implications on the projections for future prevalence. *Circulation*. 2006;114:119–125. doi: 10.1161/CIRCULATIONAHA.105.595140.
- Marini C, De Santis F, Sacco S, Russo T, Olivieri L, Totaro R, Carolei A. Contribution of atrial fibrillation to incidence and outcome of ischemic stroke: results from a population-based study. *Stroke*. 2005;36:1115–1119. doi: 10.1161/01.STR.0000166053.83476.4a.
- Hylek EM, Go AS, Chang Y, Jensvold NG, Henault LE, Selby JV, Singer DE. Effect of intensity of oral anticoagulation on stroke severity and mortality in atrial fibrillation. *N Engl J Med*. 2003;349:1019–1026. doi: 10.1056/NEJMoa022913.
- Conen D, Chae CU, Glynn RJ, Tedrow UB, Everett BM, Buring JE, Albert CM. Risk of death and cardiovascular events in initially healthy women with new-onset atrial fibrillation. *JAMA*. 2011;305:2080–2087. doi: 10.1001/jama.2011.659.
- Wolf PA, Mitchell JB, Baker CS, Kannel WB, D'Agostino RB. Impact of atrial fibrillation on mortality, stroke, and medical costs. *Arch Intern Med*. 1998;158:229–234.
- Hart RG, Pearce LA, Aguilar MI. Meta-analysis: antithrombotic therapy to prevent stroke in patients who have nonvalvular atrial fibrillation. *Ann Intern Med*. 2007;146:857–867.
- Ruff CT, Giugliano RP, Braunwald E, Hoffman EB, Deenadayalu N, Ezekowitz MD, Camm AJ, Weitz JI, Lewis BS, Parkhomenko A, Yamashita T, Antman EM. Comparison of the efficacy and safety of new oral anticoagulants with warfarin in patients with atrial fibrillation: a meta-analysis of randomised trials. *Lancet*. 2014;383:955–962. doi: 10.1016/S0140-6736(13)62343-0.
- Gladstone DJ, Bui E, Fang J, Laupacis A, Lindsay MP, Tu JV, Silver FL, Kapral MK. Potentially preventable strokes in high-risk patients with atrial fibrillation who are not adequately anticoagulated. *Stroke*. 2009;40:235–240. doi: 10.1161/STROKEAHA.108.516344.
- Camm AJ, Lip GY, De Caterina R, Savelieva I, Atar D, Hohnloser SH, Hindricks G, Kirchhof P; ESC Committee for Practice Guidelines (CPG). 2012 Focused update of the ESC guidelines for the management of atrial fibrillation: an update of the 2010 ESC guidelines for the management of atrial fibrillation: developed with the special contribution of the European Heart Rhythm Association. *Eur Heart J*. 2012;33:2719–2747. doi: 10.1093/eurheartj/ehs253.
- Fuster V, Ryden LE, Cannom DS, Crijns HJ, Curtis AB, Ellenbogen KA, Halperin JL, Le Heuzey JY, Kay GN, Lowe JE, Olsson SB, Prystowsky EN, Tamargo JL, Wann S, Smith SC Jr, Jacobs AK, Adams CD, Anderson JL, Antman EM, Hunt SA, Nishimura R, Ornato JP, Page RL, Riegel B, Priori SG, Blanc JJ, Budaj A, Camm AJ, Dean V, Deckers JW, Despres C, Dickstein K, Lekakis J, McGregor K, Metra M, Morais J, Osterspey A, Zamorano JL. ACC/AHA/ESC 2006 guidelines for the management of patients with atrial fibrillation. *Circulation*. 2006;114:e257–e354.
- Skanes AC, Healey JS, Cairns JA, Dorian P, Gillis AM, McMurtry MS, Mitchell LB, Verma A, Nattel S; Canadian Cardiovascular Society Atrial Fibrillation Guidelines Committee. Focused 2012 update of the Canadian Cardiovascular Society atrial fibrillation guidelines: recommendations for stroke prevention and rate/rhythm control. *Can J Cardiol*. 2012;28:125–136. doi: 10.1016/j.cjca.2012.01.021.
- Turakhia MP, Ullal AJ, Hoang DD, Than CT, Miller JD, Friday KJ, Perez MV, Freeman JV, Wang PJ, Heidenreich PA. Feasibility of extended ambulatory electrocardiogram monitoring to identify silent atrial fibrillation in high-risk patients: the Screening Study for Undiagnosed Atrial Fibrillation (STUDY-AF). *Clin Cardiol*. 2015;38:285–292. doi: 10.1002/clc.22387.
- Lowres N, Neubeck L, Salkeld G, Krass I, McLachlan AJ, Redfern J, Bennett AA, Briffa T, Bauman A, Martinez C, Wallenhorst C, Lau JK, Brieger DB, Sy RW, Freedman SB. Feasibility and cost-effectiveness of stroke prevention through community screening for atrial fibrillation using iPhone ECG in pharmacies: the SEARCH-AF study. *Thromb Haemost*. 2014;111:1167–1176. doi: 10.1160/TH14-03-0231.
- Claes N, Van Laethem C, Goethals M, Goethals P, Mairesse G, Schwagten B, Nuyens D, Schrooten W, Vijgen J. Prevalence of atrial fibrillation in adults participating in a large-scale voluntary screening programme in Belgium. *Acta Cardiol*. 2012;67:273–278.
- Lowres N, Neubeck L, Redfern J, Freedman SB. Screening to identify unknown atrial fibrillation: a systematic review. *Thromb Haemost*. 2013;110:213–222. doi: 10.1160/TH13-02-0165.
- Svensberg E, Engdahl J, Al-Khalili F, Friberg L, Frykman V, Rosenqvist M. Mass screening for untreated atrial fibrillation: the STROKESTOP study. 2015;131:2176–2184. doi: 10.1161/CIRCULATIONAHA.114.014343.
- Engdahl J, Andersson L, Mirskaya M, Rosenqvist M. Stepwise screening of atrial fibrillation in a 75-year-old population: implications for stroke prevention. *Circulation*. 2013;127:930–937. doi: 10.1161/CIRCULATIONAHA.112.126656.
- Gladstone DJ, Spring M, Dorian P, Panzov V, Thorpe KE, Hall J, Vaid H, O'Donnell M, Laupacis A, Côté R, Sharma M, Blakely JA, Shuaib A, Hachinski V, Coutts SB, Sahlas DJ, Teal P, Yip S, Spence JD, Buck B, Verreault S, Casaubon LK, Penn A, Selchen D, Jin A, Howse D, Mehdiratta M, Boyle K, Aviv R, Kapral MK, Mamdani M; EMBRACE Investigators and Coordinators. Atrial fibrillation in patients with cryptogenic stroke. *N Engl J Med*. 2014;370:2467–2477. doi: 10.1056/NEJMoa1311376.
- Healey JS, Connolly SJ, Gold MR, Israel CW, Van Gelder IC, Capucci A, Lau CP, Fain E, Yang S, Bailleul C, Morillo CA, Carlson M, Themeles E, Kaufman ES, Hohnloser SH; ASSERT Investigators. Subclinical atrial fibrillation and the risk of stroke. *N Engl J Med*. 2012;366:120–129. doi: 10.1056/NEJMoa1105575.
- Sandhu RK, Bakal JA, Ezekowitz JA, McAlister FA. Risk stratification schemes, anticoagulation use and outcomes: the risk-treatment paradox in patients with newly diagnosed non-valvular atrial fibrillation. *Heart*. 2011;97:2046–2050. doi: 10.1136/heartjnl-2011-300901.
- Ogilvie IM, Newton N, Welner SA, Cowell W, Lip GY. Underuse of oral anticoagulants in atrial fibrillation: a systematic review. *Am J Med*. 2010;123:638–645.

KEY WORDS: Editorials ■ atrial fibrillation

## Are We Ready for Mass Screening to Detect Atrial Fibrillation?

Jeffrey S. Healey and Roopinder K. Sandhu

*Circulation*. 2015;131:2167-2168

doi: 10.1161/CIRCULATIONAHA.115.017288

*Circulation* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2015 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7322. Online ISSN: 1524-4539

The online version of this article, along with updated information and services, is located on the  
World Wide Web at:

<http://circ.ahajournals.org/content/131/25/2167>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Circulation* is online at:  
<http://circ.ahajournals.org/subscriptions/>